

Local perceptions, uses and conservation status of *Detarium senegalense* in the Dahomey Gap (West Africa)

Patrice TREKPO

Forestry Research Laboratory, Faculty of Sciences,
University of Lomé, Togo

Biah IBITOYE

Laboratory of Applied Ecology, Faculty of Agricultural
Sciences, University of Abomey-Calavi, Cotonou, Benin

Dramane ISSIAKO

National University of Sciences, Technologies, Engineering
and Mathematics, Abomey, Benin

Aristide Cossi ADOMOU

Plant Biology Laboratory, Faculty of Science and
Technology, University of Abomey-Calavi, Cotonou, Benin

Kouami KOKOU

Forestry Research Laboratory, Faculty of Sciences,
University of Lomé, Togo

Several species, including *Detarium senegalense*, face exploitation without regard for conservation or regeneration status, posing a severe threat and contributing to their decline. Few studies addressed conservation strategies of *Detarium senegalense*. This paper aimed to document the indigenous knowledge of local communities regarding *D. senegalense* and its conservation status in the Dahomey Gap region. Data were collected through semi-structured individual interviews and focus groups involving 404 participants from thirty-eight villages, representing four phytogeographic zones in the Dahomey Gap. Collected data included local names, recognition criteria, abundance measures, and uses of *D. senegalense*. This data supported calculations of knowledge diversity, evenness index, organ usage frequency, and vulnerability indices. The Student Newman and Keuls test were used for multiple comparisons and Correspondence Analysis examined the relationships between variables. Analysis revealed that most respondents were familiar with the species (75.0%), but only half possessed knowledge of its uses. The diversity of knowledge was unevenly distributed among the respondents (IE <0.5). Significant variations in knowledge were observed among phytogeographic zones and ethnic groups, while no substantial differences were found based on gender or age. The study identified the almond as the primary organ of interest for local populations due to its involvement in international trade. The timber obtained from the tree was also used. Natural habitats of *D. senegalense* were found to be in decline, with logging being identified as the main cause of habitat destruction according to 58.5% of respondents. Consequently, the vulnerability index indicated that *D. senegalense* is highly vulnerable in its natural environment with vulnerability indices higher than 2.5 in all phytodistricts. Considering the nutritional and medicinal value of *D. senegalense*, we emphasize the need to investigate structural, morphological, and molecular aspects for conservation and sustainable use.

Keywords: *Detarium senegalense*, endogenous knowledge, Vulnerability indices, Natural habitat, Plant organs

INTRODUCTION

Natural ecosystems are indispensable suppliers of resources that serve to diverse human needs, including sustenance, artisanal materials, medicinal remedies, and spiritual significance (Lykke et al., 2004; Ipbes, 2019). Among these ecosystems, forests play a crucial role in conserving biodiversity by providing habitats for numerous animal and plant species. Moreover, they represent

a significant reservoir of natural resources for many tropical countries, making substantial contributions to biodiversity and ecological balance (N'Guessan et al., 2019). In the context of food resources, edible woody species refer to trees and shrubs that produce leaves, flowers, fruits, seeds, or other parts suitable for human consumption (Ouédraogo, 2010).

In sub-Saharan Africa, in addition to agriculture, livestock rearing, and fishing, the collection of non-timber forest products (NTFPs) stands out as a significant source of income, food, and medicinal resources for rural communities (Ezebilo and Mattsson, 2010; Mawunu et al., 2017; Tchat-chambe et al., 2017a and b; Mawunu et al., 2019). In West Africa, phylogenetic resources particularly fruit trees, play a vital role in ensuring food security and the population health (Diop, 2013; Sourou et al., 2016). Various parts or organs of many woody plants are extensively used in traditional medicine and daily practices, including timber for construction and service purposes (Fandohan et al., 2008). Wild fruits are often consumed directly at the collection or harvesting sites, while others are commercially marketed, providing income for households, and contributing to food security and improved health and nutrition of the population. The significant contribution of fruit trees to these aspects is attributed to their rich diversity (Bowe and Haq, 2010).

While forest biodiversity holds crucial importance for African rural populations, several forest species are threatened in their natural habitats due to human activities and climate variations. The potential disappearance of numerous woody species is imminent, driven not only by over-exploitation but also by improper harvesting practices and inadequate regeneration (Betti, 2001; Ouédraogo, 2008; Traoré, 2008; Agali, 2009; Belem, 2009). Many rural households rely on non-timber forest products to fulfill their nutritional, health, and construction needs (Heubach et al., 2011; Shackleton et al., 2015). According to Traoré et al. (2011), these species, which are beneficial to rural communities, face threats in their natural habitats due to deforestation, vegetation fires, extensive cattle grazing, charcoal production, abundant use of firewood, slash-and-burn cultivation, illegal logging, and uncontrolled movement of genetic material for plantation establishment. Consequently, there is an urgent necessity to conserve agroforestry species in general (Boffa, 2000; Belem, 2008; Devineau et al., 2009), with particular attention to edible species.

Enhancing our comprehension of the motivations behind the use of these plants is crucial for capturing and safeguarding local knowledge associated with indigenous species. This knowledge is vital for conservation efforts and the formulation of effective domestication strategies (Chekole et al., 2015). The integration of indigenous knowledge in the development of management and conservation strategies for native plant resources is essential to ensure their effectiveness (Avocèvou, 2011). This underscores the need for a deeper understanding of the capacity of available natural resources and their usage by the population.

Several studies have highlighted the importance of forest products, particularly the diverse services they provide to communities living near forests (Larwanou et al., 2010; Lougbégnon et al., 2011; Thiombiano et al., 2012; Yameogo et al., 2013; Houèthégnon et al., 2013; Dangbo et al., 2019; Dossa et al. 2019). According to Masengo et al. (2021 a and b), the sustainability of phylogenetic resource management depends on considering the social, cultural, and economic values held by local communities. However, several species are coveted and exploited without regard for their conservation status and, especially, their regenerative capacity, posing a serious threat and contributing to their decline. *Detarium senegalense* is a prominent example, widely used for various purposes such as food, medicinal applications, timber, and more, in several West African countries including Nigeria, Senegal, Togo, and Benin. In Nigeria, the flour derived from *D. senegalense* almonds is widely consumed and holds considerable importance in the food and pharmaceutical industries (Sowemimo et al., 2011; Malami et al., 2021). A decoction prepared from the bark is believed to be effective in treating conditions like haemorrhage, pneumonia, diarrhea, stomach disorders, digestive issues, and facilitating placenta expulsion after childbirth (Sowemimo et al., 2011). In Benin, this species faces numerous threats to its survival in the short, medium, and long term, including its use as timber wood and the existence of almond trade routes to Nigeria, Ghana, and Togo (Neuenschwander et al., 2011). In addition, Atato et al. (2010) and Dangbo et al. (2019a)

highlighted the substantial involvement of *D. senegalense* seeds in commercial trade in Togo. In this context, the survival of this species now relies on appropriate exploitation approaches coupled with techniques for natural and/or assisted regeneration while considering its vulnerability.

The vulnerability of a plant species refers to its susceptibility to the risks of decline or extinction due to inadequate harvesting methods, increasing human pressures, and climate variability within its habitat (Traoré et al., 2011). In the context of conserving this highly important forest species, it is crucial to assess its vulnerability, considering anthropogenic factors. To achieve this, it is essential to obtain reliable data on the diverse uses of the species and the level of threats across different socio-cultural groups in Benin and Togo within the Dahomey Gap.

Hence, the present study had as objective to document the endogenous knowledge of local communities regarding *D. senegalense* and its conservation status in the study area.

MATERIAL AND METHODS

Study area

The study was conducted among communities living near forested areas, whether classified or not, where *D. senegalense* is found. According to Adomou et al. (2005), Akoegninou et al. (2006), Atato et al. (2010), and Dangbo et al. (2019b), *D. senegalense* is present in the Guinean, Guineo-Sudanian, Sudanian-Guinean, and Sudanian zones. Therefore, the specific focus was on the phytodistricts of Borgou Sud and Bassila in Benin, as well as zones III and IV in Togo, all located within the Dahomey Gap in West Africa. An exploratory survey was conducted in the study area to identify potential communities for the main surveys. As a result, four municipalities (in Benin) and four prefectures (in Togo) were selected, namely Tchaourou, Ouèssè, Bantè, and Bassila in Benin, and Tchamba, Est-Mono, Akebou, and Wawa in Togo (Figure 1). In total, thirty-eight villages and hamlets scattered across all eight administrative units (municipalities or prefectures) were surveyed.

Economically, agriculture is the main activity among the communities in the study area. The crops grown include maize, yam, rice, cowpea, Bambara groundnut, and others. Perennial plantations such as cashew, coffee, cocoa, teak, and gmelina are also established. Livestock rearing, predominantly practiced by the Fulani and agro-pastoralists, is of the sedentary or transhumance husbandry type. Feeding and watering of the livestock mainly rely on natural grazing, typically along watercourses and ponds. Two types of hunting are practiced: traditional hunting with rudimentary weapons during the dry season and illegal hunting using sophisticated or crude firearms. Vegetation fires are used both in soil preparation for agriculture and in hunting activities. Fishing, on the other hand, takes place in the major watercourses during the dry season. Trade appears to be the main activity for women, constituting a significant source of income, particularly in the sale of staple foods, forest products such as firewood, charcoal, and NTFPs (fruits, seeds/nuts, or other plant parts). The communities consist of several ethnolinguistic groups and practice diverse religions, including Animism, Islam, and Christianity.

Sampling and data collect

The survey was conducted during July, November, and December 2020, as well as January and February 2021. The selection of survey locations was based on an exploratory study carried out in the areas where *D. senegalense* is known to occur, as reported in previous studies conducted in Benin and Togo (Adomou et al., 2005; Akoegninou et al., 2006; Atato et al., 2010; Dangbo et al. 2019b). The exploration aimed to study the distribution and relative abundance of the species in each phytogeographic zone of Benin and Togo. Additionally, it sought to identify the localities closest to the natural habitats in each zone where the populations have an interest in *D. senegalense*. During these preliminary surveys in the study area, it was observed that on average,

21.4% of the respondents had knowledge of the uses and habitat of the species. Thus, the sample size for the study was determined using the normal approximation of the binomial distribution proposed by (Dagnelie, 1998; Levy et Lemeshow, 2008).

With N the sample size, π (21.4%) the proportion of individuals who know and/or have knowledge of at least one use of the species, $U_{21-\alpha/2}$ the value of the standard normal variable for a risk α ($\alpha = 0.05$; $U_{21-\alpha/2} = 1.96$), and δ the margin of error set at 4%. The sample size N was calculated to be 403.8 respondents, rounded up to 404 respondents.

Each respondent, as well as focus groups, participated in a semi-structured individual questionnaire. The questionnaire encompassed the following sections: (i) gathering socio-demographic information of the respondents, including name, age, gender, ethnic group, and phytogeographic zone, (ii) exploring the diverse uses of different parts of the species, (iii) identifying criteria for distinguishing individuals of the species, and (iv) investigating the various pressures affecting the species and its habitat.

Individual and group surveys, typically consisting of 2 to 5 participants, were conducted. The use of focus groups facilitated the collection of reliable information, as it encouraged discussions and interactions among the group members, resulting in data that was collectively endorsed (Assogbadjo, 2008). In addition, observations and field verification phase was carried out. This phase involved locating the natural habitats of *D. senegalense* in the study area, which served two main purposes: firstly, to corroborate the field observations with the interview data, and secondly, to identify the sampling points for the forest inventories

Data analysis

For data analysis, the respondents were grouped into different levels of stratification (phytogeographic zone, age category, ethnic groups, and gender) based on the classification proposed by Houehanou et al. (2008) and Amouzoun et al. (2019). To explore the variations in this variable based on socio-demographic characteristics of the respondents, a generalized linear model of the Poisson family was applied.

To document local perceptions about the uses and dynamics of *D. senegalense* populations, several analytical approaches were employed. Firstly, a dynamic cross-tabulation table of different variables was created to compare various uses of *D. senegalense* organs and identify the most used ones. To assess species uses knowledge of respondents, a response rate was calculated to evaluate the level of knowledge and use of *D. senegalense* organs among different ethnic groups surveyed.

The Student Newman and Keuls test were used for multiple comparisons of knowledge levels. Additionally, an Analysis of Correspondences was conducted to examine the relationships between these organs and different ethnic groups. Ethnobotanical indices were calculated to determine the distribution of knowledge within the surveyed ethnic groups (Table 1). Furthermore, the diversity of uses for each *D. senegalense* organ were determined based on the diversity and evenness index (Table 2).

To investigate the conservation status, particularly the level of degradation of natural habitats, the threats facing the species, and its vulnerability across different socio-cultural groups, the following ethnobotanical parameters were computed. These parameters include the frequency of use and the vulnerability index of the species (Table 1). Additionally, the proportion of responses per type of habitat degradation factor was calculated. All analyses were performed using R software, version 4.2.3.

RESULTS

Ethnobotanical perceptions and uses of *D. senegalense*

Socio-demographic characteristics of respondents

In relation to the phytogeographic zones, Borgou Sud stood out as the most prevalent, representing 35.2% of the respondents, followed by Zone III at 24.1%. As for ethnicity, the Nago and related groups constitute the majority, accounting for nearly 50% of the surveyed population, while the Akposso come in at a distant second with 9.36%. In terms of age categories, adults between 31 and 59 years old make up over two-thirds of the respondents. Lastly, regarding gender, the proportions are almost equal, with approximately 50% of the respondents being female and 50% being male (Table 3).

Nomenclature

D. senegalense, also known as “arbre à suif, grand détar” in French and “dattock tree” in English, has different local names depending on the sociolinguistic groups present in the Dahomey Gap. These names are sometimes made up of two words. Examples include “N’Ténékpé N’toudja or N’tutèdiakpé” among the Anii, “Ôlô odo” among the Nagot (Table 4).

Knowledge about the species according to sociolinguistic groups

Across the entire study area, an average of 75% of the respondents were familiar with *D. senegalense*, either the tree itself or one of its organs and/or its natural habitat. However, only 50.2% were aware of at least one use of the species. Analysis indicated a significant difference in knowledge between phyto-districts and between ethnic groups. However, knowledge did not differ significantly by gender or age (Table 5). The number of uses was highest in phyto-districts IV (0.93 ± 0.31) and lowest in the Borgou Sud phytodistricts (0.03 ± 0.18). Regarding usages of the species by ethnic groups, the number of uses was highest among Moba, Ditamari, Akebou, Kabiè and Akposso, with frequencies ranging from 0.90 to 1.17. However, Fon, Peulhs, Bariba and Nagot recorded a low level of use, with frequencies lower than 0.48.

Knowledge about *D. senegalense* exhibited greater diversity in Zone IV and among the Kabiè and Moba communities (a minority) within the study area, primarily because most respondents were non-natives. Conversely, knowledge was comparatively lower in Borgou Sud phyto-district and among the Fon and associated groups (Figure 2).

Diversity and evenness indices

The study computed and presented the Shannon-Weaver diversity index (ID) and evenness index (IE) based on different factors such as phyto-districts and sociocultural layers (Table 6). The diversity index (ID) ranged from 1.61 to 23.3, with low evenness index (IE) values below 0.5, indicating diverse and heterogeneous knowledge among the respondents. Regarding gender, the knowledge diversity among women was relatively more homogeneous compared to men. Similarly, different ethnic groups show varying knowledge diversity, with all groups displaying heterogeneity. Among age groups, adults demonstrate more homogeneous knowledge diversity than the youth and the elderly.

When considering the entire study area regardless factors or categories, the evenness index remained low (0.323), indicating that the distribution of knowledge diversity among the respondents were not uniform.

Socioeconomic perceptions of *D. senegalense*

The findings revealed that local trade of *D. senegalense* almonds was common. In the phyto-district of Bassila, a significant number of women (58%) recognized the existence of a well-organised

market for *D. senegalense* almonds, while this percentage was lower (ranging from 17% to 24%) in the other three zones studied. As for sector animation, the presence of some individual intermediaries was approved by both men and women in varying proportions (42% and 85%, respectively). In Benin, the sale channel of seeds involved traders of both genders in equal percentages (100% each) in Bassila, but women (72.2%) were the primary sellers in Borgou Sud. In Togo, it was unanimously recognized that men from zone III and women from zone IV participated in the seed trade (Table 7).

Conservation status and vulnerability of *D. senegalense*

Frequency of use of *D. senegalense*

Several ways of using *D. senegalense* were observed, leading to populations using various plant parts, but in different proportions. Frequencies of use (FU) were determined for each plant parts (Figure 3). The most used part of *D. senegalense* was its seed or almond, followed by the wood. However, other parts, including fruit (pulp), leaf, bark, or root, were not used very extensively. This observation was in line with the on-site findings, where both men and women were primarily involved in activities related to fruit gathering for almond extraction and the use of wood for various purposes associated with *D. senegalense*.

Drivers of degradation of natural habitats of *D. senegalense*

The natural environments where *D. senegalense* was found correspond primarily to wet areas along watercourses, semi-deciduous dense forests, and foothills of mountains, regardless of the ethnic background of those surveyed. Most of the respondents who were acquainted with this species agreed that the existing individuals of *D. senegalense* were mostly mature specimens. This observation raised concerns about the natural regeneration of the species, highlighting the necessity of implementing measures to ensure its sustainability. Certain trees are spared during deforestation, especially in specific areas like the Akposso plateau, among the Akebou people, and the Ana (related to Nago) community. However, some of these preserved trees were unfortunately exploited for timber and charcoal production.

The investigations carried out among the local populations living near potential sites where *D. senegalense* is or was present revealed a range of perceptions regarding the level of habitat degradation, varying from low to very high (Figure 4). However, it is noteworthy that regardless of the phyto-district, most respondents (ranging from 51.3% to 74.1%) expressed concerns about a very high level of habitat degradation. This concern was particularly prominent in Borgou Sud phyto-district, where over 64% of men and 74% of women held the opinion of a significantly increasing level of degradation. Furthermore, the farmers who had *D. senegalense* trees in their garden or field claimed that they obtained them naturally and not through planting.

Almost all men and women, regardless of age or ethnicity, stated that currently, to locate *D. senegalense* trees, one must venture deeper into protected forests or other designated areas in the phyto-districts of Bassila, Borgou Sud, and Zone III. This suggests that the natural habitats and *D. senegalense* individuals were experiencing various factors leading to their destruction. However, in Zone IV, *D. senegalense* could be found in coffee, cocoa, and kola plantations, as well as in home gardens.

This habitat loss of *D. senegalense* could be explained by several drivers with various level (Table 8). In Borgou Sud, Bassila and zone III, most of respondents identified forest products exploitation (illegal and abusive logged for timber and charcoal) as the main anthropogenic drivers of habitat degradation of the species. This phenomenon was cited with high frequency in Borgou Sud phyto-district (approximately 90% of both men and women). Nevertheless, agriculture was identified as the main factor of degradation of the natural habitats of the species in Zone IV, followed by deforestation caused by logging activities. Additionally, in other phyto-districts, agriculture was

acknowledged as a secondary cause of habitat degradation for the species. The study has revealed that among respondents in the Bassila phyto-district, approximately 17% attributed negative impacts on the species' habitats to fruit gathering. However, in all four phytogeographic zones, lower than 10% of respondents assumed other activities that might negatively influence the habitats. These activities included vegetation fires, grazing, and climate change (Table 8). Interestingly, these findings align well with observations in the field. Indeed, most of *D. senegalense* trees were found in protected areas within the Bassila, southern Borgou, and Zone III phyto-districts where agricultural activities were prohibited, providing a more favorable environment for the species. In contrast, Zone IV, where agriculture is conducted without significant constraints, houses *D. senegalense* trees within open vegetation formations, making them more vulnerable to habitat degradation.

Vulnerability of *D. senegalense* in the different phyto-districts

Table 9 presents the vulnerability index of assessed parameters. The average vulnerability index of *D. senegalense* across the four zones was 2.58. It was 2.56 for the phyto-districts Borgou Sud and Bassila, then for zone III, and 2.67 for zone IV. These values are closer to 3 and indicate that the species is vulnerable in all studied phyto-districts.

DISCUSSION

Ethnobotanical and ethno-ecological knowledge of *D. senegalense* in the Dahomey Gap

Most respondents indicated that the species inhabits areas gallery forests, moist semi-deciduous forests, foot of hills and valleys. This perception was evident in certain local names for the species, particularly among the Nagots and related communities, where it is referred to as "Ôlô Odo, meaning "millstone" or "hollow" of the river. This traditional knowledge aligns with the field observations made during verifications and georeferencing of the species occurrences, and with the findings of previous researchers such as Arbonnier (2019), Adomou (2005), Akoègninou et al. (2006), Dangbo et al. (2019), Dossa et al. (2019), and Houénon et al. (2021).

In this study, the diversity of knowledge was heterogeneous, indicating that species is used for multiple purposes within each ethnic group. However, this diversity becomes more consistent when considering gender and age, with values of evenness index (IE) like those found for *Chrysophyllum albidum* and *Mansonia altissima* by Lougbégnon et al. (2016) and Wédjangnon et al. (2016), respectively (0.40 and 0.11). The values obtained in the present study are lower than those obtained by Masengo et al. (2021a) for *Lippia multiflora* (IE = 0.98). This result confirmed some previous studies that have shown that ethnobotanical knowledge was globally influenced by gender (Lougbégnon et al., 2011), educational level (Dassou et al., 2015) and profession of respondents (Beltràn-Rodríguez et al., 2014). Thus, knowledge about the species' usage was not evenly distributed among the participants. These findings are consistent with previous studies that reported several uses of *D. senegalense* in Togo (Dangbo et al., 2019a) and in Benin (Dossa et al., 2020a). For instance, *D. senegalense*, as a wild species, has specific parts, such as almonds obtained from crushing ripe fruits, which are central to commercial activities. The documented ethnobotanical knowledge about the species was closely in line with the results found by Dangbo et al. (2019a) in Togo, where over 90% of the population reported having a good understanding of the species, especially concerning the usage of almonds. Other uses were identified, notably, wood for timber, leaves, roots, and bark, corroborating the findings of Dangbo et al. (2019a), Dossa et al. (2020) and Houénon et al. (2021). The almond trade was the most important use category, followed by timber, but the food and ritual categories are virtually absent. Like Houénon et al. (2021), our results showed that *D. senegalense* fruits were not usually consumed. This finding may be linked to the fact that the fruits of this species were not considered in the eating habits of local populations.

Several studies (Amaku et al., 2021; Olayiwola, 2021; Olatunji et al, 2021; Margaret 2021) have

confirmed various physical, biochemical, pharmacological, and medicinal properties of some main parts of *D. senegalense*. Research conducted by Malami et al. (2020) in Nigeria identified *D. senegalense* as one of the medicinal plants used for cancer treatment. Moreover, Sowemimo et al. (2011) reported that the leaves are consumed as vegetables and are also employed in treating conjunctivitis. The leaf decoction is used to relieve itching and treat skin infections, as mentioned by the same authors. Additionally, the seeds are used as thickeners in specific culinary preparations. These results indicate that despite the increasing demand for *D. senegalense* in the sub-region, only a limited number of culinary and medicinal applications are currently made from its organs.

Regarding the dynamics of *D. senegalense* populations, it was noted that the observation was consistent across phytodistricts. The low abundance of *D. senegalense* population was mentioned by most respondents. This finding confirmed the results of Dangbo et al. (2019b), Dossa et al. (2019), and Houénon et al. (2021), who also reported low individual density of *D. senegalense* in their respective study areas. The excessive and systematic harvesting of fruits for almond commercialisation was considered one of the main underlying causes of this situation (Adomou et al., 2011; Dangbo et al., 2019a; Dossa et al., 2020b; Houénon et al., 2021). In Togo, the sale of almonds contributes to the income of local populations (Dangbo et al., 2019a). Additionally, there was intense cutting-down of adult individuals of the species. There is a trade of *D. senegalense* almonds in zone IV. The commercial name given to the seed is “bethor”. The profit generated by this commerce for the collectors was low due to the informal nature of the entire commercialisation chain, despite the desire of communities to see this trade well-organized and structured to ensure better valorisation and conservation of the species. These results were not in line with those of Diop (2008), who reported significant income derived from the ditax trade both for collectors and the state. This author stated that *D. senegalense* holds a prominent position in the list of important forest fruit species in the Senegalese economy.

The almonds of *D. senegalense* were not yet directly consumed by local communities but were subject to commercial transactions. These results corroborated those of Dangbo et al. (2019a). Regarding the actors involved in the collection of *D. senegalense* fruits, the present results revealed that both women and men participate in this activity. These findings were in line with Atato et al. (2010), who concluded that *D. senegalense* seeds were collectively harvested, with even men participating. This highlighted the importance of women in the management of forest resources. Despite their participation in the exploitation of forest resources, they are often overlooked in the forest policy development. This results in a limited consideration of their needs and strategic interests during the implementation of management plans (Bauer, 2010). However, the lack of gender consideration represents a threat for resources.

Conservation status and vulnerability of *D. senegalense* in Dahomey Gap

This study revealed that the existing populations of *D. senegalense* consist mainly of older specimens, and natural regeneration is minimal or virtually absent. The natural habitats of the species are undergoing significant decline due to deforestation and agricultural activities. These findings are consistent with previous research by Houénon et al. (2021) and Sambiéni et al. (2015), which also reported declining populations and high levels of degradation of forest ecosystems in Benin. Similarly, other studies in West Africa have shown a general trend of vegetation regression (Adjonou et al., 2010; Houéssou et al., 2012; BNTD, 2016; Bouko et al., 2016; Adingra, 2017; Silue et al., 2023). This concordance of the perception of the populations in the study area with the results of previous studies demonstrated the relevance of exploring endogenous knowledge about the spatio-temporal dynamic of natural habitats of *D. senegalense*. Indeed, several authors have reported that, in addition to the scientific understanding of past vegetation, it is crucial to consider the insights of local communities. These insights help identify any change in the floral landscape (Ali et al., 2023; Hahn-Hadjali and Thiombiano, 2000). Similarly, in the context of managing natural resources holistically, Busquet (2006) highlighted the critical value of considering the traditional ecological knowledge of local communities. The author also pointed out that human perceptions

significantly influence the definition of conservation policies, as conservation practices often align with prevailing social values.

The vulnerability of a species is influenced by various factors, including the usage, the specific organ harvested, the frequency of use, the collection method, the abundance of the species, and the stage of development (Traoré et al., 2011; Ayéna et al., 2016; Nzuki, 2016; Ngbolua, 2020). Additionally, the species morphological type (Dassou et al., 2014), the biotope (Masengo et al., 2022), and its natural regenerative capability play significant roles. This vulnerability increases when the harvested plant organ struggles to regenerate, or its collection leads to the systematic cutting-down of the entire tree (Betti, 2001). According to Guigma et al. (2012), the survival of a species is adversely affected by every exploited organ. The vulnerability index values observed in different studied phytogeographic zones were in line with the findings of Agbo et al. (2017) and Masengo et al. (2022). These results can be attributed to the fact that for *D. senegalense*, significant organs are obtained from large trees, particularly in the case of fruit and almond collection, which is usually gathered from the ground rather than being plucked from the tree. As mentioned above, the main threats to species lie in the cutting of trees and the limited natural regeneration, which firmly establishes its vulnerability. In the Dahomey Gap, the exploitation, transformation, and trading of NTFPs offer opportunities to address food shortages and generate substantial financial gains during periods of scarcity, as demonstrated by studies of Mawunu et al. (2019; 2020), and Ngbolua et al. (2019b; 2021). In addition to its very low capacity for natural regeneration, the mode of exploitation and social conditions of communities of West Africa make *D. senegalense* highly vulnerable in the Dahomey gap. Given that *D. senegalense* does not yet possess a notable significant economic value like the baobab (*Adansonia digitata*), the African locust bean (*Parkia biglobosa*), shea tree (*Vitellaria paradoxa*), or butter tree (*Pentadesma butyracea*), for instance, there is still an opportunity for future research to emphasize the improvement techniques for its regeneration and domestication. It is imperative to rely on the conviction of local populations who have lived for decades with natural resources and confronted with perpetual environmental changes because they are the first to identify solutions and strategies for better adaptation (Ali et al., 2023).

CONCLUSION

The study has revealed that *D. senegalense*, a well-known indigenous plant resource, facing several threats making it vulnerable in Dahomey Gap. Local knowledge about the species varies across different phyto-districts and ethnic groups. However, the diversity of knowledge was evenly distributed based on gender and age. The main use of the species was the collection, processing, and commercialisation of its almonds. Logging and agriculture posed significant threats to the natural habitat of *D. senegalense*, making it highly vulnerable in the studied areas. To preserve the species, the study recommended supporting natural regeneration and exploring domestication strategies. Conservation efforts should involve local communities and their traditional knowledge. Collaboration is essential in developing comprehensive conservation plans that consider the vulnerability of *D. senegalense* and its sustainable use.

REFERENCES

- Adingra M.A.O. (2017). Dynamique du peuplement et stock de carbone dans la mosaïque de végétation de la forêt classée de Bamo (Côte d'Ivoire). PhD thesis, University of Félix Houphouët-Boigny, Côte d'Ivoire, 158 p.
- Adjonou K., Djiwa O., Kombate Y., Kokutse A.D., Kokou K. (2010). Étude de la dynamique spatiale et structure des forêts denses sèches reliques du Togo: implications pour une gestion durable des aires protégées. *Int. J. Biol. Chem. Sci.*, 4: 168-183.
- Adjossou K. (2004). Diversité floristique des forêts riveraines de la zone écologique iv du Togo. DEA

de biologie de développement option: biologie végétale appliquée, Université de Lomé, 99p.

Adomou C.A. (2005). Vegetation Patterns and Environmental gradients in Benin Implications for biogeography and conservation. PhD Thesis Wageningen University, Wageningen, The Netherlands 133p.

Adomou C.A, Agbani O.P., Sinsin B. (2011). Plants In: Neuenschwander P, Sinsin B, Goergen G (eds) Protection de la nature en Afrique de l'Ouest: Une Liste Rouge pour le Bénin Nature Conservation in West Africa: Red List for Benin. International Institute of Tropical Agriculture, pp 21-46.

Agbo R.I., Missihoun A.A., Vihotogbe R., Assogbadjo E.A., Ahanhanzo C., Agbangla C. (2017). Impacts des usages traditionnels sur la vulnérabilité de *Detarium microcarpum* Guill & Perr (Caesalpiniaceae) dans le district phytogéographique Zou au Bénin (en Afrique de l'Ouest). , 11: 730-742.

Akoègninou A., Van der Burg W.J., Van der Maesen L.J.G. (2006). Flore analytique du Bénin. Backhuys Publishers, Leiden, Netherlands 1034 pp.

Ali M., Amadou Issoufou A., Abdramane S., Soumana I., Mahamane A. (2023). Importances des services écosystémiques dans la résilience des populations rurales dans le Sud-Ouest du Niger: synthèse bibliographique. *Int. J. Biol. Chem. Sci.*, 17: 2076-2088.

Amouzoun B.M.A.A, Badoun B.R., Ahamide D.Y.I., Adomou C.A. (2019). Connaissances ethnobotaniques et conservation de *Uvariadendron angustifolium* (Engl & Diels) R E Fries (Annonaceae) dans l'îlot forestier Ewe- Ada-Kplame au Sud-Est de Bénin, Afrique de l'Ouest. *Rev. Ivoir. Sci. Technol.*, 34: 328-348.

Amaku J.J.F., Ngwu C.M., Ogundare S.A., Akpomie K.G., Edozie O.I., Conradie J. (2021). Thermodynamics, kinetics and isothermal studies of chromium (VI) biosorption onto *Detarium senegalense* stem bark extract coated shale and the regeneration potentials. *Int. J. Phytoremediation*, 23:1-11.

Arbonnier M. (2019). Arbres, arbustes et lianes de zones sèches d'Afrique de l'Ouest. 2e éd, Cirad Mnhn, Paris, France, 541 p.

Assogbadjo A.E., Glèlè Kakai R., Chadaré F.J., Thomson L., Kyndt T., Sinsin B., Van Damme P. (2008). Folk classification, perception, and preferences of baobab products in West Africa: consequences for species conservation and improvement. *Eco. Bot.*, 62: 74-84.

Atato A., Wala K., Batawila K., Woegan A.Y., Akpagana K. (2010). Diversité des fruitiers spontanés du Togo. *Fruit, Veg. Cereal Sci. Biotech.*, 4:1-9.

Avocèvou-Ayisso C.A. (2011). Étude de la viabilité des populations de *Pentadesma butyracea* Sabine et de leur socio-économie au Bénin. PhD thesis in Agricultural sciences, University of Abomey-Calavi, 223 p.

Ayéna A.C., Agassounon D., Tchiboza M., Assogbadjo A.E., Adoukonou-Sagbadja H., Mensah G.A., Agbangla C., Ahanhanzo C. (2016). Usages et vulnérabilité de *Pterocarpus santalinoides* L'her ex de (Papillionoidae), une plante utilisée dans le traitement des gastro-enterites dans le Sud du Bénin. *Eur. Sci. J.*, 12:218-231

Bauer L. (2010). Forêts et réduction de la pauvreté dans les pays en développement: une relation à déchiffrer. Essai de master en ingénierie et management de l'environnement et du développement durable, Université de Sherbrooke, p119.

- Belem B., Smith O.C., Theilade I., Bellefontaine R., Guinko S., Mette Lykke A., Diallo A., Boussim J.I. (2008). Identification des arbres hors forêt préférés des populations du Sanmatenga (Burkina Faso). *Bois For. Trop.*, 298: 53-64.
- Beltrán-Rodríguez L., Ortiz-Sánchez A., Mariano N.A., Maldonado-Almanza B., Reyes-García V. (2014). Factors affecting ethnobotanical knowledge in a mestizo community of the Sierra de Huautla Biosphere Reserve. Mexico. *J. Ethnobiol. Ethnomed.*, 10: 2-18.
- Betti J.L. (2001). Vulnérabilité des plantes utilisées comme antipaludiques dans l'arrondissement de Mintom au sud de la réserve de Biosphère du Dja (Cameroun). *Syst. & Geogr. Pl.*, 71: 661- 678.
- BNETD (2015). Rapport pour les états généraux de la forêt, de la faune et des ressources en eau: gestion durable des ressources forestières, Côte d'Ivoire. 117p
- Boffa J.M. (2000). Les parcs agroforestiers en Afrique subsaharienne. *Cahier FAO Conservation*, 34, 258 p.
- Bousquet F., Castella J.C., Ekasingh B., Hoanh C.T., Kam S.P., Manichon H., Ni D.V., Quang D.D., Trébuil G., Tuong T.P. (2006). Recherche éco-régionale sur la gestion intégrée des ressources naturelles dans les écosystèmes de riziculture en Asie du sud-est IN: La France et le CGIAR: des résultats scientifiques pour la recherche agricole internationale. Colombo, Sri Lanka: CGIAR Challenge Program on Water and Food: 85-90.
- Bowe C., Haq N. (2010). Quantifying the global environmental niche of an underutilised tropical fruit tree (*Tamarindus indica*) using herbarium records. *Agric. Ecosyst. Environ.*, 139: 51-58.
- Byg A., Balsvel H. (2001). Diversity and use of pâlms in Zahamena, eastern Madagascar. *Biodivers. Conserv.*, 10: 951-970.
- Cavin A.L. (2007). Contribution à la connaissance taxonomique et chimique de fruits africains du genre «*Detarium*» (Fabaceae - Caesalpinioideae): «*D microcarpum*» Guill et Perr et des formes comestibles et toxiques de «*D senegalense*» JF Gmel. PhD thesis: Univ Genève, no Sc 3838.
- Chatelain C., Kadjo B., KoneI Refisch J. (2001). Relation Faune - Flore dans le Parc National de Tai: une étude bibliographique Tropembos-Cote d'Ivoire Série 36345481.
- Chekole, G., Asfaw, Z., Kelbessa, E. (2015). Ethnobotanical study of medicinal plants in the environs of Tara-gedam and Amba remnant forests of Libo Kemkem District, northwest Ethiopia. *J Ethnobiology Ethnomedicine*, 11:4, 38p.
- Dagnelie P. (1998). *Statistique théorique et appliquée*. vol 2, Paris, De Boeck et Larcier, Belgique.
- Dangbo F.A., Adjonou K., Kokou K., Blaser J. (2019a). The socio-economic contribution of *Detarium senegalense* seeds to rural livelihoods in Togo (West Africa). *Int. J. Biol. Chem. Sci.*, 13: 1582-1595.
- Dangbo F.A., Adjonou K., Kokou K., Blaser J. (2019b). Ecological Sustainability of Detar (*Detarium senegalense* J F Gmel) in Togo (West Africa). *Am. J. Plant Sci.*, 10: 417-432.
- Dassou H.G., Ogni C.A., Yédomonhan H., Adomou A.C., Tossou M., Dougnon J.T., Akoègninou A. (2014). Diversité, usages vétérinaires et vulnérabilité des plantes médicinales au Nord-Bénin. *Int. J. Biol. Chem. Sci.*, 8: 189-210.
- Dassou H.G., Yedomonhan H., Adomou A.C., Ogni C.A., Tossou M.G., Akoegninou A. (2015). Facteurs socioculturels et environnementaux déterminant la connaissance ethno-vétérinaire au Bénin. *Afrique Science*, 11: 335-360.



Devineau J.L., Fournier A., Nignan S. (2009). Ordinary biodiversity" in western Burkina Faso (West Africa): what vegetation do the state forests conserve? *Biodivers Conservation*, 18: 2075-2099.

Diop N. (2008). Collecte de Ditax dans les zones inventoriées au niveau de la région de Fatick Amélioration de la compétitivité des boissons et nectars à base de fruits locaux au Sénégal. Proj FNRAA-ITA, Rapp Mission, Dakar.

Dossa A.K.B., Ouinsavi C., Houetchegnon T.N., Sourou N.B. (2019). Caractérisation structurale des peuplements naturels de *Detarium senegalense* JF Gmel (Caesalpiniaceae) au Bénin, *Science de la vie, de la terre et agronomie, Rev. Ramres.*, 7:15-23.

Dossa B.A.K., Ouinsavi C., Houetchegnon T., Sourou B.N. (2020a). Knowledge points and research perspectives on *Detarium Senegalense*, a vulnerable species in Benin. *Int. J. Res. Stud. Biosci.*, 8: 4-12.

Dossa B.A.K., Ouinsavi C., Houetchegnon T., Sourou B.N. (2020b). Germination of Seeds and Growth in the Nursery and Field of Seedlings of *Detarium senegalense* in Benin. *Eur. Sci. J.*, 16: 12.

Ezebilo E.E., Mattsson L. (2010). Contribution of non-timber forest products to livelihoods of communities in southeast Nigeria. *Int. J. Sust. Dev. World Ecol.*, 17:231-235.

Fandohan B., Glèlè Kakai R., Sinsin B., Pelz D. (2008). Caractérisation dendrométrique et spatiale de trois essences ligneuses médicinales dans la forêt Classée de Wari-Marou au Bénin. *Rev. Ivoir. Sci. Technol.*, 12: 173-186.

Guigma Y., Zerbo P., Millogo-Rasolodimby J. (2012). Utilisation des espèces spontanées dans trois villages contigus du Sud du Burkina Faso. *Tropicultura*, 30: 230-235.

Hahn-Hadjali K., Thiombiano A. (2000). Perception des espèces en voie de disparition en milieu Gourmantché (Est du Burkina Faso). *Berichte des Sonderforschungsbereichs*, 268: 285-297.

Heubach K., Wiitg R., Nuppenau E., Hahn K. (2011). The economic importance of nontimber forest products (NTFPs) to livelihood maintenance of rural west African communities: a case study from northern Benin. *Ecol Econ.*,70:1991-2001.

Houéhanou D.T., Assogbadjo A.E., Chadare F.J., Zanzo S., Sinsin B. (2016). Approches méthodologiques synthétisées des études d'ethnobotanique quantitative en milieu tropical. *Annales des Sciences Agronomie*, 20: 187-205.

Houénon G.H.A., Djossou A.K., Kouhinkpo E.Y., Salako K.V., Tchobo F.P., Adomou A.C., Yédomonhan H. (2021). Parataxonomy, perceived dynamics and diversity of uses of two *Detarium* species in Benin (West Africa). *Genetic Resources and Crop Evolution*, 68:1-28.

Houéssou G.L., Loughbegnon O.T., Gbesso G.H.F., Anagonou S.E.L., Sinsin B. (2012). Ethnobotanical study of the African star apple (*Chrysophyllum albidum* G Don) in the Southern Benin (West Africa). *J. ethnobiol. ethnomed.*, 8: 1-10.

Houèchégnon T., Gbèmavo D.S.J.C., Ouinsavi C., Sokpon N. (2015). Ethnobotanical knowledge and traditional management of african mesquite (*Prosopis africana*) populations in Benin, West Africa. *Journal of Ethno. and Tradi. Med. Photon*, 125: 1124-1135.

Ilomuanya M.O., Ekerebe Z. Cardoso-Daodu I., Sowemimo A. (2020). Formulation and Evaluation of Sunscreen Cream Using *Detarium senegalense* Oil as Base. *Tropical Journal Natural Product Research*, 4:141-145.

- IPBES (2019). Communiqué de presse: le dangereux déclin de la nature: un taux d'extinction des espèces «sans précédent» et qui s'accélère Paris, France.
- Larwanou M., Oumarou I., Snook L., Danguimbo I., Eyog-Matig O. (2010). Pratiques sylvicoles et culturelles dans les parcs agroforestiers suivant un gradient pluviométrique nord-sud dans la région de Maradi au Niger. *Tropicultura*, 28: 115-122.
- Levy P.S., Lemeshow S. (2008). *Sampling of populations: Methods and Applications*. New York, USA, Wiley, 192 p.
- Lougbégnon T.O., Tente B.A.H., Amontcha M., Codjia J.T.C. (2011). Importance culturelle et valeur d'usage des ressources végétales de la réserve forestière marécageuse de la vallée de Sitatunga et zones connexes *Bulletin de la Recherche Agronomique du Bénin*, 70.
- Lougbégnon O.T., Nassi K.M., Gbesso G.H.F. (2015). Ethno-botanique quantitative de l'usage de *Chrysophyllum albidum* G Don par les populations locales au Bénin. *Journal of Applied Biosciences*, 95: 9028-9038.
- Lykke A.M., Kristensen M.K., Ganaba S. (2004). Valuation of local use and dynamics of 56 woody species in the Sahel. *Biodiversity Conservation*, 13: 1961-1990.
- Malami I., Jagaba N.M., Abubakar I.B., Muhammad A., Alhassan A.M., Peter M.W., Yahaya P.M., Mshelia I.Z., Mathias H.E.S.N. (2020). Integration of medicinal plants into the traditional system of medicine for the treatment of cancer in Sokoto State, Nigeria. *Heliyon Journal*, 6: e04830.
- Masengo C.A., Djolu R.D., Ngbolua K.N., Nagahuedi J.M.S., Ilumbe B.I., Mpiana P.T. (2022). Étude ethnobotanique et vulnérabilité de *Vitex doniana* Sweet (Lamiaceae) dans la forêt péri-urbaine de Gbado-Lite, RDC. *Rev. Mar. Sci. Agron. Vét.*, 10: 179-184.
- Masengo C.A., Bongo N.G., Robijaona B., Ilumbe G.B., Ngbolua K.N., Mpiana, P.T. (2021a). Étude ethnobotanique quantitative et valeur socioculturelle de *Lippia multiflora* à Kinshasa, République Démocratique du Congo. *Rev. Mar. Sci. Agron. Vét.*, 9: 93-101.
- Masengo C.A., Inkoto C.L., Munsebi J.M., Mandjo B.L., Mpiana P.T., Ngbolua K.N. (2021b). Connaissance et usages de *Quassia africana* par les peuples Mongo, Yaka et Yombe de Kinshasa en République Démocratique du Congo. *Rev. Mar. Sci. Agron. Vét.*, 9: 737-745.
- Mawunu M., Dionisio C.A., Lukoki L., Ngbolua K.N., Luyindula N. (2019). Ethnobotanical and Socio-economics of *Dracaena camerooniana* Baker in Uíge Province, Northern Angola. *J. Agric. Ecol.*, 20: 1-15 .
- Mawunu M., Eduardo A.S.O., Balomba P., Mohindo A.A., Bongo G., Ngbolua K.N. (2017). Food Security and Livelihood of Rural Households of Songololo Territory in Kongo Central Province, Democratic Republic of the Congo. *International Journal of Health Economics and Policy*, 2: 97-103.
- Mawunu M., Pedro P., Lautenschläger T., Biduayi F.M., Kapepula P.M., Ngbolua K.N., Luyeye F.L., Luyi N. (2020). Nutritional Value of Two Underutilized Wild Plant Leaves Consumed as Food in Northern Angola: *Mondia whitei* and *Pyrenacantha klaineana*. *European J. Nutr. Food Saf.*, 12: 116-127.
- N'Guessan A.E., Kassi N.J., Yao N.O., Amani H.K.B., Gouli G.Z.R., Pioniot C., Irie Bi C.Z., Hérault B. (2019). Drivers of biomass recovery in a secondary forested landscape of West Africa. *Forest Ecology and Manag.*, 433: 325-331.
- Neuenschwander P., Sinsin B., Goergen G. (eds). (2011). *Protection de la Nature en Afrique de*

l'Ouest: Une Liste Rouge pour le Bénin Nature Conservation in West Africa: Red List for Benin. International Institute of Tropical Agriculture, Ibadan, Nigeria 365 p.

Ngbolua K.N., Nzamonga G.A., Gbatea K.A., Nzale M.S., Masengo A.C., Ndolete G.J.P., Bongo N.G., Zakwani L.N., Libwa M.T.B., Yangba T.S., Gerengbo K.G. (2019b). Knowledge on Non-Timber Forest Products (NTFPs) Marketed in Democratic Republic of the Congo: A Case Study of Gbadolite City and Surroundings, Nord Ubangi. *Agricultural and Biological Sciences Journal*, 5: 20-28.

Ngbolua K.N. (2020). *Ethnobotanique quantitative: Approches méthodologiques pour l'évaluation et la valorisation du savoir endogène en régions tropicales*. Éditions Universitaires Européennes, Riga: Latvia.

Ngbolua K.N., Molongo M.M., Libwa M.T.B., Amogu J.J.D., Kutshi N.N, A Masengo C.A. (2021). Enquête ethnobotanique sur les plantes sauvages alimentaires dans le Territoire de Mobayi-Mbongo (Nord-Ubangi) en République démocratique du Congo. *Rev. Mar. Sci. Agron. Vét.*, 9: 259-265.

Nzuki B.F. (2016). *Recherches ethnobotaniques sur les plantes médicinales dans la Région de Mbanza-Ngungu, RDC*. PhD thesis (PhD), Faculté des Sciences en Bio-Ingénierie, Université de Gand, Belgique, 349 p.

Okafor J.C. (1981). Essences forestières pour agroforesterie dans le Sud du Nigeria In «MAC Donald L H1981, Agroforesterie en Afrique tropicale humide». *Compte rendu d'un colloque tenu à Ibadan, Nigéria 27 avril-1er mai 1981NRST-17F/UNUP-467188p*.

Olayiwola Y.B. (2020). Effect of solvents on *Detarium senegalense* bark and characterization of the resulting bio-oil for pf resins system. *Pro Ligno*, 16:19-26.

Ouédraogo I. (2010). *Fonctions et services des phytocénoses de la réserve partielle de faune de Pama-nord et des zones riveraines*. Mémoire de DEA en sciences biologiques appliquées Université de Ouagadougou, Burkina Faso, 50p.

Ouédraogo I. (2008). *Diversité des espèces ligneuses utiles de la région du Nord du Burkina Faso; état des peuplements de cinq espèces d'importance socio-économique*. Mémoire d'Ingénieur en Eaux et Forêts Université de Bobo-D, Burkina Faso, 68p.

Sambiéni K.R., Mama A., Toyi M.S. (2015). Perception paysanne sur la fragmentation du paysage de la Forêt classée de l'Ouémé Supérieur au nord du Bénin. *VertigO - la revue électronique en sciences de l'environnement*, 15: 1-18.

Sambuc M.C. (1887). *Contribution à la flore et à la matière médicale de la Sénégambie*. Ec Sup Pharm, Thèse, Montpellier, France, pp 84-94.

Shackleton C.M., Pandey A.K., Ticktin T. (2015). *Ecological sustainability for Non-timber Forest Products: Dynamics and Case Studies of Harvesting*. Routledge, New York, USA, 280p.

Silué A.P., Kouassi K. E., Soro D., Kouakou A.M.M., Soro D. (2023). Dynamique paysagère et particularités floristiques de la forêt classée de Pouniakélé (Nord-Ouest de la Côte d'Ivoire). *Rev. Mar. Sci. Agron. Vét.*, 11: 185-192.

Sourou B.N., Yabi J., Ouinsavi C., Sokpon N. (2016). Importance socio-économique de la prune rouge (*Haematostaphis barteri* Hook F) au Bénin. *Int. J. Biol. Chem. Sci.*, 10: 326-343.

Sowemimo A.A., Pendota C., Okoh B., Omotosho T., Idika N., Adekunle A.A., Afolayan A.J. (2011). Chemical composition, antimicrobial activity, proximate analysis and mineral content of the seed of



Detarium senegalense JF Gmelin. African Journal of Biotechnology, 10: 9875-9879.

Tchatchambe N.B.J., Solomo E.B., Kirongozi B.F., Lebisabo B.C., Dhed'a D.B., Tchatchambe W.B.J., Ngombe K.N., Mpiana P.T., Mbemba F.T., Ngbolua K.N. (2017a). Évaluation de la valeur nutritive et des facteurs antinutritionnels de quatre légumes alimentaires sauvages consommées à Kisangani et ses environs (Province de la Tshopo, RD Congo). Int. J. of Innov. and Sci. Res., 30: 75-90.

Tchatchambe N.B.J., Solomo E.B., Kirongozi B.F., Lebisabo B.C., Dhed'a D.B., Tchatchambe W.B.J., Ngombe K.N., Mpiana P.T., Mbemba F.T., Ngbolua K.N. (2017b). Analyses nutritionnelle et toxicologique de trois plantes alimentaires traditionnelles de la Tshopo en République Démocratique du Congo. Int. J. of Innov. and Sci. Res., 30: 105-118.

Thiombiano D.N.E., Lamien N., Dibong D.D., Boussim I.J., Belem B. (2012). Le rôle des espèces ligneuses dans la gestion de la soudure alimentaire au Burkina Faso. Séch., 23: 86-93.

Traoré L., Ouédraogo I., Ouédraogo A., Thiombiano A. (2011). Perceptions, usages, et vulnérabilité des ressources végétales ligneuses dans le Sud-Ouest du Burkina- Faso. Int. J. Biol. Chem. Sci., 5: 258.

Wédjangnon A.A., Houèthégnon T., Ouinsavi, C. (2016). Caractéristiques ethnobotaniques et importance socio-culturelle de *Mansonia altissima* A Chev in Bénin, Afrique de l'Ouest. Journal of Animal & Plant Sciences., 29: 4678-4690.

Yameogo G., Yelemou B., Boussim I.J., Traoré D. (2013). Gestion du parc agroforestier du terroir de Vipalogo (Burkina Faso): Contribution des ligneux à la satisfaction des besoins des populations. International Journal of Biology and Chemical Sciences 7: 1087-1105.

References