

# Evaluation of the antioxidant and antibacterial activities of Tunisian *Artemisia Herba-alba* essential oil

Ameni DHIFALLAH

Hamadi ROUISSI

Houcine SELMI

Sylvo-Pastoral Institute of Tabarka, University of Jendouba, Tunisia

Ecole Supérieure d'Agriculture de Mateur, Tunisia

Sylvo-Pastoral Institute of Tabarka, University of Jendouba, Tunisia

Tunisia is characterized by a climate that allows the proliferation of many plants rich in active substances with multiple biological activities and can replace the use of antioxidants and synthetic antibiotics. For this reason, *Artemisia* leaves and twigs were collected from the central region of Tunisia (Thala). The essential oil was extracted using hydro-distillation and analyzed using GC/MS, the antioxidant activity of *Artemisia Herba-alba* was evaluated by the DPPH test and the antibacterial powers against four bacterial strains was measured by the agar well method diffusion. GC/MS results showed that the main components of *Artemisia Herba-alba* essential oil were  $\alpha$ -thujone (23.9 %) and chrysanthenone (17.4%). Indeed, the results showed a potent antioxidant effect (85.2 % inhibition of free radicals DPPH) and the IC<sub>50</sub> value was 84.8  $\mu$ g/ml. Concerning the antibacterial activity, the oil was active against Gram-negative and Gram-positive bacteria. A strong effect was observed against *Salmonella* (29mm) and *Bacillus* (22.5 mm). To conclude, the antioxidant power and the antibacterial activity are strongly correlated with the chemical composition of the essential oil.

**Keywords:** *Artemisia Herba-alba*, essential oil, antioxidant activity, antibacterial activity

## Introduction

During the last decades, a concern about the undesirable effects of synthetic molecules intended for the fight against oxidative stress and bacterial infections. It therefore seems important to find an alternative to the use of synthetic antioxidants and conventional antibiotics (Rosenthal et al., 2012). Medicinal plants are considered as an essential raw material source for the discovery of new molecules necessary for the development of future drugs, without side effects (Benbelaïd et al., 2014). An example of these metabolites are essential oils and plant extracts (Rahman and Sattar, 2011). The genus *Artemisia* belongs to the Asteraceae family, *Artemisia* species are widely used in traditional medicine. Over 300 species of this genus are found mainly in arid and semi-arid areas of Europe, America, North Africa and Asia (Nikolova et al., 2010). The development of new antioxidants and antibiotics of better quality and of lower toxicity is essential. For this purpose, the investigation of plants represents an invaluable potential (Adams and Gmünder, 2007).

In this context, this study aims to evaluate the antioxidant activity of *Artemisia Herba alba* the essential oil collected from the region of Thala (Tunisia) by the reducing power and the inhibition of the 2,2-diphenyl-1-picrylhydrazyl radical (DPPH), as well as its antibacterial activity against gram positive and gram negative strains.

## Material and methods

### Plant material

The aerial part (Leaf, Stem) of *Artemisia Herba-alba* were collected from the region of Thala (Tunisia) (latitude 35° 35.583' N, longitude 008° 39.717' E, altitude 847 m, semi-arid bioclimate).

### **Essential oil extraction**

The extraction of essential oils was carried out by steam distillation, this method consists in placing the plant material on a grid located a few centimeters from the bottom of the extractor filled with water. The steam enriched with volatile constituents is condensed under the effect of a cooling system. The floral water is collected in a glass balloon and the separated essential oil is collected in an opaque glass bottle. The extraction was carried out for four hours at a temperature of 100 °C.

### **Essential oil gas chromatography analysis**

The essential oil obtained is diluted with n-pentane (Hosni et al., 2008), then analyzed by high resolution gas chromatography using an Agilent 6890 brand device (Agilent Technologies España, SL, Las Rozas, Spain) equipped with a flame ionization detector (FID), an auto-injector, an autosampler and an Rtx-1 non-polar column (30 m\* 0.32 mm, film thickness 0.25 µm). The system is controlled by GC-Solution type software which ensures the electronic integration of the different peaks.

### **Identification of volatile compounds of essential oil**

The volatile compounds of essential oils were identified by calculating their retention index (IR) from a range of linear alkanes (C8-C25) injected under the same analytical conditions (Van Den Dool and Kratz, 1963). The calculation of the retention indices for volatile compounds is given by the following equation:

$$RI = [n + (TR_i - TR_n) / (TR_{(n+1)} - TR_n)] * 100$$

RI: unidentified peak retention index, n: number of carbon atoms of the aliphatic hydrocarbon eluted just before the peak to identify,  $TR_i$ : retention time of the unidentified peak,  $TR_n$ : retention time of the aliphatic hydrocarbon eluted just before the peak to be identified,  $TR_{n+1}$ : retention time of the aliphatic hydrocarbon eluted just after the peak to be identified

### **1, 1-Diphenyl-2-picrylhydrazyl radical (DPPH) scavenging activity**

The antioxidant capacity of *Artemisia Herba Alba* essential oil was evaluated according to the method described by (Braca et al., 2002). The diluted essential oil solutions were prepared in methanol. Ascorbic acid was used as the standard. Then 1 mL of 0.004% DPPH solution was mixed with 1 mL of sample solution, the mixture was then shaken and kept in the dark for 30 min and optical density was measured at 517 nm. The radical-scavenging activities, expressed as percentage inhibition of DPPH, were calculated according to the following equation:

$$I (\%) = [(A_0 - A_1) / A_0] * 100$$

I: DPPH inhibition (%),  $A_0$ : absorbance of the blank,  $A_1$ : absorbance of the extract/standard.

The concentration of sample required for 50% inhibition was determined and represented as  $IC_{50}$  for each of test solution, which is expressed as µg/ml. All measurements were performed in triplicate.

### **Antibacterial activity**

The antibacterial activity of the essential oil was measured by means of the agar well diffusion test described by Güven et al. (2006) against four bacterial strains (*Listeria monocytogenes* (foodstuff

2132), *Bacillus subtilis* ATCC 6633, *Escherichia coli* ATCC 25922, *Salmonella enterica* (foodstuff). A suspension (100  $\mu$ l) of bacteria was spread on the plates of nutrient agar, then 6 mm wells were bored using a sterile cork borer. 60  $\mu$ l of essential oil were placed into the wells. Finally, they were incubated at 37°C for 48 h and the antibacterial activity was evaluated by measuring zone of inhibition. The tests were carried out in triplicate.

### Statistical analysis

All data were subjected to statistical analysis by the variance according to the GLM procedure of the software SAS 2002 and compared by Duncan multiple rank test (Duncan, 1955). The model used was:

$$Y_{ij} = \mu + A_i + E_{ijk}$$

Where  $Y_{ij}$ : dependent variable.  $\mu$ : overall of Y;  $A_i$ : effect of the  $i$ th essential oil;  $E_{ijk}$ : residual error.

## Results and discussion

### Chemical composition

30 compounds of *Artemisia Herba alba* essential oil are identified and listed in table 1, representing 99.5% of the oil. The chemical class distributions of the essential oil could be separated into five classes, Monoterpenes represented about 94.9 % of the total constituents of essential oils. Oxygenated monoterpenes constituted the main chemical class of the oil (74.2 %) and they were represented by 9 derivatives, with  $\beta$ -thujone (23.9 %), chrysanthenone (17.4 %),  $\alpha$ -thujone (10.3 %) and 1,8-cineole (9.78 %) as main components. In addition, monoterpene hydrocarbons were represented by 11 compounds representing 20.7 % of all oil. Among these compounds, p-cymene (8.27 %),  $\alpha$ -pinene (3.25 %) and Sabinene (3.31 %) were the most important. Sesquiterpenes constituted 4.6 % of all oil, among them oxygenated derivatives (3 compounds) represented only 1.5% while hydrocarbons (7 compounds) reached 3.1% of all oil.

The chemical variability of *Artemisia Herba alba* observed in the Tunisian areas (north-west, center and south-east of Tunisia) showed various compositions dominated either by a single component ( $\alpha$ -thujone, camphor, chrysanthenone or trans-sabinyacetate) or characterized by the presence of two or more of these compounds at appreciable levels (Boukrich et al., 2010). Indeed, in our study thujones ( $\alpha$ -thujone and  $\beta$ - thujone) present the highest percentage 34.26% of all oil, as reported by Haouari and Ferchichi (2009) thujone are the components which give the specific odor and taste to *Artemisia Herba-alba* plants.

For further comparison, Chrysanthenone reached a high percentage (17.4%), according to (Neffati et al., 2008; Akrouf, 2004) this component was not found in the Tunisian *Artemisia Herba alba* essential oil at such high concentrations.

### Antioxidant activity of *Artemisia Herba-alba* essential oil

The results reported in Figure 1 showed that for each concentration, the antioxidant activity of *Artemisia Herba alba* is lower than ascorbic acid which is known by its high antioxidant potential. At a concentration of about 300  $\mu$ g/mL, the exhibited greatest inhibitory activity reaches as high as 85.2 %. The IC<sub>50</sub> value obtained for scavenging activity on DPPH radical were evaluated (IC<sub>50</sub>=84.8  $\mu$ g/ml) and showed that the oil possessed a potent antioxidant effect (Table 2). From the results reported in Table 1, the property of *Artemisia Herba alba* of being a radical scavenging agent was attributed to its richness in oxygenated monoterpenes (Kadri et al., 2011) or to the synergistic effect of more than one oil compound (Younsi et al., 2016).

Disagreeing with our current data, (Akrouit et al.,2010)reported that the essential oil of leaves of *Artemisia Herba-alba* growing in the wild in southern Tunisia exhibited low anti-free radical activity in the method (DPPH). However, the same result was found for different essential oils of aerial parts of *Artemisia Herba-alba* cultivated in southern Tunisia (Mighri et al., 2010).

### **Antibacterial activity of *Artemisia Herba-alba* essential oil**

The in vitro antibacterial activity of *Artemisia Herba Alba* was qualitatively and quantitatively assessed by the diameter of inhibition zone, MIC and MBC values.

The data indicated in table 3 that the oil was active against Gram negative bacteria and Gram-positive bacteria. An important effect was noticed against *Salmonella enterica* and *Bacillus subtilis* ATCC 6633 which were very sensitive with an inhibition diameter of 29 and 22.5 mm respectively. Furthermore, *Escherichia coli* ATCC 25922 and *Listeria monocytogenes* were tightly less sensitive with an inhibition zone of 21.3 and 18 mm.

The results in table 4 show that the oils exhibited varying levels of antibacterial activity against the investigated bacteria ( $p < 0.05$ ). all bacteria are inhibited at concentrations ranging from 10  $\mu\text{L/mL}$  to 100  $\mu\text{L/mL}$ . Those concentrations are higher than the doses found by Sbayou et al.(2014) and Mighri et al.(2010). In fact, our essential oil showed a significant bacteriostatic activity as specified by the ratio MBC/MIC of 5 to 10 that is  $>4$  (Pankey et al., 2004). This activity could be related to the amounts of oxygenated mono-and sesquiterpene hydrocarbons (Rahman and Sattar, 2011). The presence of  $\beta$ -thujone as a major compound in *Artemisia Herba alba* may be responsible for its antibacterial activity. Thus, it is difficult to attribute the activity of a complex mixture to a single or particular constituent. Major or trace compounds might give rise to the antimicrobial activity exhibited. Possible synergistic and antagonistic effect of compounds in the oil should also be taken into consideration (Sbayou et al., 2014).

## **Conclusion**

The present work has shown that the essential oil of *Artemisia Herba alba* collected from the center of Tunisia is doubted of considerable antioxidant and antibacterial activities, hence the possibility of using them as antioxidants and natural antibiotics. This biological activity is strongly linked to the chemical composition of the essential oil and to the concentration of these aromatic and medicinal plant on secondary metabolites. Research on this essential oil should be continued to better estimate other potential as anti-inflammatory, anti diabetic and antifungal activity, even use it as supplement in the field of animal feeding.

## **References**

Adams M.,Gmünder F., Hamburger M.(2007). Plants Traditionally Used in Age Related Brain Disorders-A Survey of Ethnobotanical Literature. *Journal of Ethnopharmacology*, 113: 363-381

Akrouit A., El Jani H., Zammouri T., Mighri H., Neffati M.(2010). Phytochemical screening and mineral contents of annual plants growing wild in the southern of Tunisia. *Journal of Phytology*, 2:34-40.

Akrouit A.(2004). Essential oil study of some pastoral plants from Matmata (south Tunisia) (in French).*Cah. Options Med.*, 62: 289-292.

Benbelaïd F., Khadir A., Abdoune M.A., Bendahou M., Muselli A., Costa J. (2014). Antimicrobial activity of some essential oils against oral multidrug-resistant *Enterococcus faecalis* in both planktonic and biofilm state. *Asian Pacific journal of tropical biomedicine*, 4:463-472.

Boukrich F., Zouari S., Neffati M., Abdelly C., Liu K., Casanova J., Tomi F. (2010). Chemical variability of *Artemisia herba-alba* Asso growing wild in semi-arid and arid land (Tunisia). *Journal of Essential Oil Research*, 22:331-335.

Braca A., Sortino C., Politi M., Morelli I., Mendez J. (2002). Antioxidant activity of flavonoids from *Licania licaniaeflora*. *Journal of ethnopharmacology*, 79: 379-381.

Duncan D.B.(1955) Multiple range and multiple F test. *Biometrics*, 11: 1-42.

Güven K., Yücel E., Cetintaş F. (2006) Antimicrobial Activities of Fruits of *Crataegus* and *Pyrus* Species. *Pharmaceutical Biology*, 44:79-83.

Haouari M., Ferchichi A. (2009). Essential oil composition of *Artemisia herba-alba* from Southern Tunisia. *Molecules*, 14: 1585-1594.

Hosni K., Msaâda K., Taârit M.B., Ouchikh O., Kallel M., Marzouk B.(2008). Essential oil composition of *Hypericum perforatum* L. and *Hypericum tomentosum* L. growing wild in Tunisia. *Industrial crops and products*, 27:308-314.

Kadri A., Chobba I.B., Zarai Z., Békir A., Gharsallah N., Damak M., Gdoura R. (2011). Chemical constituents and antioxidant activity of the essential oil from aerial parts of *Artemisia herba-alba* grown in Tunisian semi-arid region. *African Journal of Biotechnology*, 10: 2923-2929.

Mighri H., Hajlaoui H., Akrouit A., Najjaa H., Neffati M.(2010).Antimicrobial and antioxidant activities of *Artemisia herba-alba* essential oil cultivated in Tunisian arid zone. *Comptes Rendus Chimie*,13:380-386.

Neffati A., Skandrani I., Sghaier M.B., Bouhlel I., Kilani S., Ghedira K., Neffati M., Cherif, I., Hammami M., Chekir-Ghedira L.(2008). Chemical composition, mutagenic and antimutagenic activities of essential oils from (Tunisian) *Artemisia campestris* and *Artemisia herba-alba*. *J. Essen. Oil Res.*, 20: 471-477.

Nikolova M., Gussev C.H, Nguyen T. (2010). Evaluation of the Antioxidant action and flavonoid composition of *Artemisia* species extracts. *Biotechnology & Biotechnological Equipment*, 24:101-103.

Pankey G.A., Sabath L.D. (2004). Clinical relevance of bacteriostatic versus bactericidal mechanisms of action in the treatment of Gram-positive bacterial infections. *Clinical infectious diseases*, 38:864-870.

Rahman A., Sattar M.A.(2011). Antioxidant Properties of Essential oil and Organic Extracts of *Premna integrifolia* Linn. In International Conference and Exhibition on Pharmaceutical Biotechnology. Hyderabad: Omics Group Conferences.

Rosenthal V.D., Bijie H., Maki D.G., Mehta Y., Apisarnthanarak A., Medeiros E.A., Jayatilleke K.(2012). International Nosocomial Infection Control Consortium (INICC) report, data summary of 36 countries, for 2004-2009. *American journal of infection control*, 40:396-407.

SAS User's Guide.(2002). Version 9.00 for Windows, SAS Inst. Inc., Cary, NC, USA.

Sbayou H., Ababou B., Boukachabine K., Manresa A., Zerouali K., Amghar S. (2014). Chemical composition and antibacterial activity of *Artemisia herba-alba* and *Mentha pulegium* essential oils. *Journal of Life Sciences*, 8(1).

Van Den Dool H., Kratz, P.D.A.(1963). Generalization of the retention index system including linear



temperature programmed gas-liquid partition chromatography (No. Research).

Younsi F., Trimech R., Boulila A., Ezzine O., Dhahri S., Boussaid M., Messaoud C. (2016). Essential oil and phenolic compounds of *Artemisia herba-alba* (Asso.): Composition, antioxidant, anti acetylcholinesterase, and antibacterial activities. *International journal of food properties*, 19: 1425-1438.

## References